Important Formulas

Celsius To Fahrenheit Scale

$$T_F = \frac{9}{5} T_c + 32$$
$$T_F = 1.8 T_c + 32$$

Celsius to Kelvin Scale

$$T_K = 273 + T_C$$

7.1. Normal human body temperature is $98.6^{\circ}F$. Convert it into Celsius scale and Kelvin scale. Given Data

Temperature in Fahrenheit = $T_F = 98.6$ °F

To Find

Temperature in Celsius =
$$T_C$$
 = ?
Temperature in Kelvin = T_K = ?

Solution

By using Fahrenheit scale

$$T_{F} = \frac{9}{5} T_{c} + 32$$

$$T_{F} = 1.8 T_{c} + 32$$

$$98.6 = 1.8 T_{c} + 32$$

$$98.6 - 32 = 1.8 T_{c}$$

$$66.6 = 1.8 T_{c}$$

$$\frac{66.6}{1.8} = T_{c}$$

$$37 = T_{c}$$

$$T_{c} = 37^{\circ}\text{C}$$

Now by using Kelvin scale

$$T_K = 273 + T_C$$

 $T_K = 273 + 37$
 $T_K = 310 K$

7.2. At what temperature Celsius and Fahrenheit thermometer reading would be the same? Given Data

Let temperature in Celsius = $T_C = T$ And temperature in Fahrenheit = $T_F = T$

To Find

Temperature at which = $T = T_C = T_F$

Solution

By using Fahrenheit scale

$$T_{F} = \frac{9}{5}T_{c} + 32$$

$$T = \frac{9}{5}T + 32 \qquad \because T = T_{C} = T_{R}$$

$$T = 1.8T + 32$$

$$T - 1.8T = 32$$

$$-0.8T = 32$$

$$T = \frac{32}{-0.8}$$

$$T = -40$$

OR

$$T_F = \frac{9}{5} T_c + 32$$

$$T = \frac{9}{5} T + 32 \qquad \because T = T_C = T_F$$

$$T = \frac{9 T + 160}{5}$$

$$5T = 9T + 160$$

$$5T - 9T = 160$$

$$-4T = 160$$

$$T = \frac{160}{-4}$$

$$T = -40$$

7.3. Convert 5°F to Celsius and Kelvin scale. Given Data

Temperature in Fahrenheit = $T_F = 5$ °F

To Find

Temperature in $Kelvin = T_K = ?$

Solution

By using Fahrenheit scale

$$T_{F} = 1.8 T_{c} + 32$$

$$5 = 1.8 T_{c} + 32$$

$$5 - 32 = 1.8 T_{c}$$

$$-27 = 1.8 T_{c}$$

$$\frac{-27}{1.8} = T_{c}$$

$$-15 = T_{c}$$

$$T_{c} = -15^{\circ}\text{C}$$

Now by using Kelvin scale

$$T_K = 273 + T_C$$

 $T_K = 273 + (-15)$
 $T_K = 273 - 15$
 $T_K = 258 K$

7.4. What is equivalent temperature of 25°C on Fahrenheit and Kelvin scales?

Given Data

Temperature in Celsius = T_C = 25°C

To Find

Temperature in Fahrenheit = T_F =? Temperature in Kelvin = T_K =?

Solution

By using formula of Fahrenheit scale

$$T_F = 1.8 T_c + 32$$

 $T_F = 1.8 (25) + 32$
 $T_F = 45 + 32$
 $T_F = 77 \,^{\circ}\text{F}$

Now by using Kelvin scale

$$T_K = 273 + T_C$$

 $T_K = 273 + 25$
 $T_K = 298 K$

7.5. The ice and steam points on an ungraduated thermometer are found to be $192 \, mm$ apart. What temperature will be on Celsius scale if the length of mercury thread is at $67.2 \, mm$ above the ice point mark?

Given Data

Length between ice and steam points = l_{100} = 192 mm Length of mercury thread = l_{θ} = 67.2 mm

To Find

Temperature on Celsius scale = $T_C = ?$

Solution

We use the formula for temperature on a linear scale:

$$T_C = \frac{l_\theta}{l_{100}} \times 100$$

$$T_C = \frac{67.2}{192} \times 100$$

 $T_C = 35^{\circ}$ C

7.6. The length between the fixed point of liquid-inglass thermometer is 20 cm. If the mercury level is 4.5 cm above the lower mark, what is the temperature on the Fahrenheit scale?

Given Data

A. Payyab (GHS Chriastian Daska)

To Find

Solution

We use the formula for temperature on a linear scale:

$$T_C = \frac{l_\theta}{l_{100}} \times 100$$

$$T_C = \frac{4.5}{20} \times 100$$

$$T_C = 22.5^{\circ}C$$

$$T_F = 1.8 T_c + 32$$

 $T_F = (1.8)(22.5) + 32$
 $T_F = 40.5 + 32$
 $T_{-} = 72.5$ °F

Prepared By: M. Tayyab, SSE(Math) Govt Christian High School, Daska. Mobile: 03338114798 Website: https://hira-science-academy.github.io Page 2 of 2